

The number of permutations with a given number of sequences

Marcus Kollar

*Theoretical Physics III, Center for Electronic Correlations and Magnetism,
Institute for Physics, University of Augsburg, 86135 Augsburg, Germany
Marcus.Kollar@physik.uni-augsburg.de*

October 15, 2006

$P(n, s)$ denotes the number of permutations of $1, 2, \dots, n$ that have exactly s sequences. Canfield and Wilf [math.CO/0609704] recently showed that $P(n, s)$ can be written as a sum of s polynomials in n . We determine these polynomials explicitly and also obtain explicit expressions for $P(n, s)$ and its fixed- s generating function $u_s(x)$.

1 Introduction

A permutation of the numbers $1, 2, \dots, n$ has at least two sequences, where a *sequence* or *run* is defined as the maximal interval of consecutive increasing or decreasing numbers. Let $P(n, s)$ denote the number of permutations of n numbers with s runs. Its basic recurrence is [1]

$$P(n, s) = sP(n-1, s) + 2P(n-1, s-1) + (n-s)P(n-1, s-2), \quad (n \geq 2) \quad (1a)$$

$$P(2, s) = 2\delta_{s,1}, \quad (1b)$$

Recently Canfield and Wilf [2] showed that the fixed- s generating function of $P(n, s)$ has the form

$$u_s(x) \stackrel{\text{def}}{=} \sum_{n=2}^{\infty} P(n, s)x^n = \frac{\Phi_s(x)}{\Delta_s(x)}, \quad \Delta_s(x) \stackrel{\text{def}}{=} \prod_{i=0}^{s-1} (1 - (s-i)x)^{\lfloor i/2 \rfloor + 1}, \quad (s \geq 1) \quad (2)$$

where $\Phi_s(x)$ is a polynomial of degree $1 + \lceil s(s+2)/4 \rceil$, i.e., one larger than the degree of the polynomial $\Delta_s(x)$ in the denominator. Furthermore they showed that $P(n, s)$ can be written as

$$P(n, s) = \sum_{i=0}^{s-1} \psi_i(n, s)(s-i)^n, \quad (n-1 \geq s \geq 1) \quad (3)$$

where each $\psi_i(n, s)$ is a polynomial in n of degree at most $\lfloor i/2 \rfloor$. These polynomials satisfy

$$\psi_i(n, s) = \frac{s\psi_i(n-1, s) + 2\psi_{i-1}(n-1, s-1) + (n-s)\psi_{i-2}(n-1, s-2)}{s-i}, \quad (s-1 \geq i \geq 1) \quad (4a)$$

$$\psi_{-1}(n, s) = 0, \quad \psi_0(n, s) = K(s) \stackrel{\text{def}}{=} 2^{-(s-2)}, \quad (4b)$$

so that (3) satisfies (1).

In Section 2 we determine the polynomials $\psi_i(n, s)$ from (4) and an explicit formula for $P(n, s)$ from (3). An auxiliary generating function is evaluated in Section 3. In Section 4 we obtain explicit expressions for the generating functions $u_s(x)$ and the polynomials $\Phi_s(x)$.

2 The polynomials $\psi_i(n, s)$

We put $\phi_i(n, t) \stackrel{\text{def}}{=} \psi_i(n, s - i)$. From (4) we find

$$t\phi_i(n, t) = (t + i)\phi_i(n - 1, t) + 2\phi_{i-1}(n - 1, t) + (n - t - i)\phi_{i-2}(n - 1, t), \quad (t \geq 1, i \geq 1) \quad (5a)$$

$$\phi_{-1}(n, t) = 0, \quad \phi_0(n, t) = K(t), \quad (5b)$$

which has the advantage that t is merely a parameter but not involved in the recurrence.

From (5) we find that the generating function $f(x, n, t) \stackrel{\text{def}}{=} \sum_{i=0}^{\infty} \phi_i(n, t)x^i$ satisfies

$$tf(x, n, t) = (2x + (n - 2)x^2 + (1 - x^2)t)f(x, n - 1, t) + (1 - x^2)x\frac{\partial}{\partial x}f(x, n - 1, t), \quad (6a)$$

$$f(0, n, t) = K(t). \quad (6b)$$

We seek a solution of the form $f(x, n, t) = g(x, t)h(x, t)^n$. After inserting this into (6a) we first eliminate all terms proportional to n by choosing $h(x, t) = (1 - x^2)^{1/2}$. This leads to a separable linear differential equation for $g(x, t)$, which is solved by $g(x, t) = (1 - x)^{1/2}(1 + x)^{-3/2}(1 + (1 - x^2)^{1/2})^{-t}$. Using (6b) we arrive at

$$f(x, n, t) = K(t)(1 - x)^2(1 - x^2)^{(n-3)/2} \left(\frac{2}{1 + \sqrt{1 - x^2}} \right)^t. \quad (7)$$

In order to determine the coefficient of x^i of (7) we use [3]

$$(1 - x^2)^{(n-3)/2} = \sum_{k=0}^{\infty} a_k(n)x^{2k}, \quad a_k(n) \stackrel{\text{def}}{=} \binom{(n-3)/2}{k} (-1)^k, \quad (8)$$

$$\left(\frac{2}{1 + \sqrt{1 - x^2}} \right)^t = \left(\frac{1 - \sqrt{1 - x^2}}{x^2/2} \right)^t = \sum_{m=0}^{\infty} b_m(t)x^{2m}, \quad b_m(t) \stackrel{\text{def}}{=} \frac{t(2m + t - 1)!}{m!(m + t)!4^m}. \quad (9)$$

We find

$$f(x, n, t) = K(t)(1 - x)^2 \sum_{j=0}^{\infty} p_j(n, t)x^{2j}, \quad p_j(n, t) \stackrel{\text{def}}{=} \sum_{k=0}^j a_k(n)b_{j-k}(t). \quad (10)$$

With the convention $p_{-1}(n, t) = 0$ we finally obtain

$$\phi_{2j}(n, t) = K(t)(p_j(n, t) + p_{j-1}(n, t)), \quad \phi_{2j+1}(n, t) = -2K(t)p_j(n, t), \quad (j \geq 0) \quad (11)$$

as well as

$$\psi_i(n, s) = (-1)^i K(s - i) \left(p_{\lfloor \frac{i}{2} \rfloor}(n, s - i) + p_{\lfloor \frac{i-1}{2} \rfloor}(n, s - i) \right) \quad (12)$$

$$= K(s - i) \sum_{j=0}^{\lfloor i/2 \rfloor} g_{i,j} p_j(n, s - i), \quad g_{i,j} \stackrel{\text{def}}{=} \begin{cases} \delta_{j, i/2} + \delta_{j, i/2-1} & i \text{ even} \\ -2\delta_{j, (i-1)/2} & i \text{ odd} \end{cases}. \quad (13)$$

Since $a_j(n)$ is a polynomial in n of order j , so are $p_j(n, t)$, $\phi_{2j}(n, t)$, and $\phi_{2j+1}(n, t)$. Thus $\psi_i(n, s)$ is a polynomial in n *precisely* of degree $\lfloor i/2 \rfloor$; it was already shown in Ref. [2] that this degree is *at most* $\lfloor i/2 \rfloor$. The first few $\psi_i(n, s)$ are given in Table 1.

Inserting (13) into (3) immediately yields an explicit formula for $P(n, s)$,

$$P(n, s) = \sum_{i=0}^{s-1} K(s - i)(s - i)^n \sum_{j=0}^{\lfloor i/2 \rfloor} g_{i,j} p_j(n, s - i), \quad (n - 1 \geq s \geq 1) \quad (14)$$

which will be used again in Section 4.

3 The auxiliary generating functions $A_k(z)$

Before turning to $u_s(x)$ we evaluate the auxiliary generating functions $A_k(z) \stackrel{\text{def}}{=} \sum_{n=2}^{\infty} a_k(n)z^n$. Their generating function is

$$\begin{aligned} A(y, z) &\stackrel{\text{def}}{=} \sum_{k=0}^{\infty} A_k(z)y^k = \sum_{n=2}^{\infty} (1-y)^{(n-3)/2} z^n \\ &= \frac{1}{\sqrt{1-y}} \frac{z^2}{1-z\sqrt{1-y}} = \left(z + \frac{1}{\sqrt{1-y}} \right) \frac{z^2}{1-z^2(1-y)} \\ &= \sum_{k=0}^{\infty} \left(\frac{z^{2k+3}}{(1-z^2)^{k+1}} + \sum_{m=0}^{\infty} \binom{m-1/2}{k} z^{2m+2} \right) (-y)^k. \end{aligned} \quad (15)$$

The inner sum evaluates to

$$\sum_{m=0}^{\infty} \binom{m-1/2}{k} x^m = \binom{-1/2}{k} {}_2F_1\left(\frac{1}{2}, 1; \frac{1}{2} - k; x\right) = \binom{-3/2}{k} \frac{(-x)^k}{(1-x)^{k+1}} {}_2F_1\left(-k, \frac{1}{2}; \frac{3}{2}; x^{-1}\right). \quad (16)$$

$A_k(z)$ is then obtained as

$$A_k(z) = \frac{z^2(-1)^k \tilde{A}_k(z)}{(1-z^2)^{k+1}}, \quad \tilde{A}_k(z) \stackrel{\text{def}}{=} z^{2k+1} + \binom{-3/2}{k} \sum_{m=0}^k \binom{k}{m} \frac{(-1)^{k+m}}{2m+1} z^{2k-2m}, \quad (17)$$

where $\tilde{A}_k(z)$ is a polynomial in z of degree $2k+1$. We evaluate its k th derivative at $z = -1$ and find

$$\left. \frac{d^k \tilde{A}_k(z)}{dz^k} \right|_{z=-1} = \binom{-3/2}{k} \frac{k!}{k+1} \left[-4^k + \sum_m \binom{k}{m} \binom{2k-2m}{k-2m} \frac{(-1)^m(k+1)}{2m+1} \right] = 0, \quad (18)$$

since the sum which appears inside the square brackets equals 4^k , as certified by the WZ [3] proof certificate $R(k, m) = 4m(2m+1)(2k-2m+1)(k+1-2m)^{-1}(k+1)^{-1}$. Thus $z = -1$ is an at least $(k+1)$ -fold zero of $\tilde{A}_k(z)$, i.e., the polynomial $\tilde{A}_k(z)$ contains a factor $(1+z)^{k+1}$.

For the higher derivatives of $\tilde{A}_k(z)$ at $z = -1$ we find

$$\begin{aligned} \tilde{a}_k(p) &\stackrel{\text{def}}{=} \frac{(-1)^k}{(k+p+1)!} \left. \frac{d^{k+p+1} \tilde{A}_k(z)}{dz^{k+p+1}} \right|_{z=-1} \\ &= (-1)^p \left[\binom{2k+1}{k+p+1} - \binom{-3/2}{k} \sum_{m=0}^{\lfloor (k-p-1)/2 \rfloor} \binom{k}{m} \binom{2k-2m}{k+p+1} \frac{(-1)^{k+m}}{2m+1} \right]. \quad (0 \leq p \leq k) \end{aligned} \quad (19)$$

The remaining sum (which vanishes for $p = k$) does not seem to allow further simplification.

After inserting $\tilde{A}_k(z) = (-1)^k \sum_{p=0}^k \tilde{a}_k(p)(1+z)^{k+p+1}$ into (17), the final expression for $A_k(z)$ becomes

$$A_k(z) = \frac{\tilde{\Phi}_k(z)}{(1-z)^{k+1}}, \quad \tilde{\Phi}_k(z) = z^2 \sum_{p=0}^k \tilde{a}_k(p)(1+z)^p. \quad (k \geq 0) \quad (21)$$

Note that the polynomial $\tilde{\Phi}_k(z)$ in the numerator of $A_k(z)$ has degree $k+2$, one larger than the degree of the polynomial in the denominator.

4 The generating functions $u_s(x)$ and polynomials $\Phi_s(x)$

For the generating functions $u_s(x)$ we employ (14) and $\sum_{n=2}^{\infty} p_j(n, t) z^n = \sum_{k=0}^j A_k(z) b_{j-k}(t)$, whence

$$\begin{aligned} u_s(x) &= \sum_{i=0}^{s-1} K(s-i) \sum_{j=0}^{\lfloor i/2 \rfloor} g_{i,j} \sum_{k=0}^j A_k((s-i)x) b_{j-k}(s-i), \quad (s \geq 1) \\ &= \sum_{i=0}^{s-1} \sum_{k=0}^{\lfloor i/2 \rfloor} \frac{B_{i,k}(x, s-i)}{(1-(s-i)x)^{k+1}}, \quad B_{i,k}(x, t) \stackrel{\text{def}}{=} K(t) \tilde{\Phi}_k(tx) \sum_{j=k}^{\lfloor i/2 \rfloor} g_{i,j} b_{j-k}(t). \end{aligned} \quad (22)$$

Clearing denominators yields $u_s(x) = \Phi_s(x)/\Delta_s(x)$ as in (2), with

$$\Phi_s(x) = \sum_{i=0}^{s-1} \left(\prod_{\substack{m=0 \\ m \neq i}}^{s-1} (1-(s-i)x)^{\lfloor m/2 \rfloor + 1} \right) \sum_{k=0}^{\lfloor i/2 \rfloor} B_{i,k}(x, s-i) (1-(s-i)x)^{\lfloor i/2 \rfloor - k}.$$

The polynomial $\Phi_s(x)$ indeed has degree $1 + \lceil s(s+2)/4 \rceil$, as proven already in [2]. The first few $\Phi_s(x)$ are given in Table 2.

5 Conclusion

We derived explicit expressions for $P(n, s)$ and $u_s(x)$ using the formulation of Ref. [2] in terms of polynomials $\psi_i(n, s)$ and $\Phi_s(x)$. Previously, Carlitz derived explicit formulas for $P(n, s)$ based on a two-variable generating function [4, 5, 6]. However, Canfield and Wilf [2] note that the final formulas in Ref. [5] are not entirely correct. We checked our results by computer algebra for $i, s, n \leq 20$.

References

- [1] D. André, *Études sur les maxima, minima et séquences des permutations*, Ann. Sci. École Norm. Sup. **3** (1884), 121–135.
- [2] E. R. Canfield and H. S. Wilf, *Counting permutations by their runs up and down*, Preprint arxiv.org/math.CO/0609704 (2006).
- [3] H. S. Wilf, *generatingfunctionology* (Academic, San Diego, 1994).
- [4] L. Carlitz, *Enumeration of permutations by sequences*, Fibonacci Quart. **16** (1978), 259–268.
- [5] L. Carlitz, *The number of permutations with a given number of sequences*, Fibonacci Quart. **18** (1980), 347–352.
- [6] L. Carlitz, *Enumeration of permutations by sequences — II*, Fibonacci Quart. **19** (1981), 398–406; *ibid.*, 465.

i	$\psi_i(n, s)$
-1	0
0	$K(s)$
1	$K(s-1)(-2)$
2	$K(s-2)(-2n+s+8)/4$
3	$K(s-3)(2n-s-3)/2$
4	$K(s-4)(4n^2-4(s+8)n+s^2+15s+32)/32$
5	$K(s-5)(-8n^2+8(s+3)n-2(s^2+5s+10))/32$
6	$K(s-6)(-8n^3+12(s+8)n^2-2(3s^2+45s+98)n+s^3+21s^2+74s+144)/384$
7	$K(s-7)(16n^3-24(s+3)n^2+4(3s^2+15s+32)n-2(s^3+6s^2+23s+42))/384$
8	$K(s-8)(16n^4-32(s+8)n^3+8(3s^2+45s+100)n^2-8(s^3+21s^2+76s+160)n$ $+s^4+26s^3+107s^2+442s+768)/6144$
9	$K(s-9)(-16n^4+32(s+3)n^3-8(3s^2+15s+34)n^2+8(s+3)(s^2+3s+16)n$ $-(s^4+6s^3+35s^2+126s+216))/3072$
10	$K(s-10)(-32n^5+80(s+8)n^4-80(s^2+15s+34)n^3+40(21s^2+s^3+78s+176)n^2$ $-2(5s^4+130s^3+555s^2+2510s+4504)n$ $+s^5+30s^4+115s^3+870s^2+2824s+4800)/122880$

Table 1: The first few polynomials $\psi_i(n, s)$.

s	$\Phi_s(x)$
1	$2x^2$
2	$4x^3$
3	$2x^4(5-6x)$
4	$4x^5(24x^2-29x+8)$
5	$2x^6(720x^4-1704x^3+1436x^2-501x+61)$
6	$4x^7(17280x^6-51336x^5+61188x^4-37256x^3+12209x^2-2041x+136)$
7	$2x^8(-3628800x^9+15729120x^8-29341872x^7+30810864x^6-20028656x^5+8353808x^4$ $-2236439x^3+370871x^2-34601x+1385)$
8	$4x^9(696729600x^{12}-3555239040x^{11}+8107966944x^{10}-10906662240x^9+9627417336x^8$ $-5872225480x^7+2537780728x^6-783164808x^5+171355239x^4-25936503x^3+2579241x^2$ $-151385x+3968)$
9	$2x^{10}(1316818944000x^{16}-8712886694400x^{15}+26410986334080x^{14}-48618945021312x^{13}$ $+60779114417952x^{12}-54684478479456x^{11}+36624658707312x^{10}-18628018251952x^9$ $+7273896122392x^8-2188789058612x^7+506111568077x^6-89028957282x^5+11685816855x^4$ $-1107016832x^3+71414171x^2-2804314x+50521)$
10	$4x^{11}(2528292372480000x^{20}-18993012627456000x^{19}+66507291476582400x^{18}$ $-144199874533248000x^{17}+216971940209451264x^{16}-240735551604776064x^{15}$ $+204330019791468672x^{14}-135856983272339904x^{13}+71875337579512880x^{12}$ $-30562090468050280x^{11}+10504633067351272x^{10}-2924633644527940x^9+658629666786430x^8$ $-119364099863329x^7+17244871619376x^6-1956223222079x^5+170214919190x^4$ $-10952481287x^3+490431140x^2-13630637x+176896)$

Table 2: The first few polynomials $\Phi_s(x)$.